



Operads, clones, and distributive laws

Pierre-Louis Curien

► To cite this version:

Pierre-Louis Curien. Operads, clones, and distributive laws. Operads and Universal Algebra: Proceedings of China-France Summer Conference, Jul 2010, Tianjin, China. pp.25-50. hal-00697065

HAL Id: hal-00697065

<https://hal.science/hal-00697065>

Submitted on 14 May 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Operads, clones, and distributive laws

Pierre-Louis Curien

(PPS Laboratory, CNRS & University Paris Diderot, Paris, France)

May 14, 2012

Abstract

We show how non-symmetric operads (or multicategories), symmetric operads, and clones, arise from three suitable monads on **Cat**, each extending to a (pseudo-)monad on the bicategory of categories and profunctors. We also explain how other previous categorical analyses of operads (via Day's tensor products, or via analytical functors) fit with the profunctor approach.

1 Introduction

Operads, in their coloured and non-symmetric version, are also known as multicategories, since they are like categories, but with morphisms which have a single target as codomain but a multiple source (more precisely, a list of sources) as domain. These morphisms are often called operations (whence the term “operad” for the whole structure). Operads for short are the special case where the multicategory has just one object, and hence the arity of a morphism is just a number n (the length of the list), while the coarity is 1. There are many variations.

- non-symmetric operads versus symmetric operads versus clones (these variations concern the way in which operations are combined to form compound operations);
- operations versus cooperations (one input, several outputs), or “bioperations” (several inputs, several outputs);
- operations whose shape, or arity, is more structured than a list (it could be a tree, etc...).

In this paper, we deal principally with the first variation, and touch on the second briefly. Our goal is to contribute to convey the idea that these variations can be smoothly and rather uniformly understood using some categorical abstractions.

We are aware of two approaches for a general categorical account of such variations. The first one, which is also the earliest one, is based on spans,


while the second one is based on profunctors. In both approaches, one abstracts the details of the variant in a monad, and one then extends the monad to the category of spans, or to the category of profunctors. Here, we take the profunctor approach.

- Spans are pairs of morphisms with the same domain, in a suitable category \mathbf{C} . Burroni has shown [[8]] that every cartesian¹ monad T on \mathbf{C} extends to a monad on spans, and then, for example, multicategories arise as endomorphisms endowed with a monoid structure in the (co)Kleisli (bi)category of a category of spans.
- Profunctors are functors $\Phi : \mathbf{C} \times \mathbf{C}'^{op} \rightarrow \mathbf{Set}$. As pointed out by Cheng [[11]], every monad in \mathbf{Cat} (the 2-category of categories) satisfying a certain distributive law can be extended to the category of profunctors (whose objects are categories and whose morphisms are profunctors). We exhibit here how

non-symmetric and symmetric operads, and clones, arise (again) as monoids in the (co)Kleisli (bi)category associated with the respective extended monad.

This is all rather “heavy” categorical vocabulary. We shall unroll this slowly in what follows. We offer the following intuitions for why monads, spans, and profunctors are relevant here.

- Monad. Consider the following composition of operations:



By plugging the output of g and h on the inputs of f in parallel, we put together the three inputs of g and the input of h , yielding a compound operation with $3 + 1 = 4$ inputs. In other words, one may read two shapes on the upper part of the picture: $((\cdot, \cdot, \cdot), \cdot)$, and $(\cdot, \cdot, \cdot, \cdot)$. The first one remembers the construction, the second one flattens it. This flattening is a typical monad multiplication (monad of powersets, of lists, etc...).

- Span. Each bioperation has an arity and a coarity: the legs of the span are this arity and coarity mappings, respectively.

¹A monad T is cartesian if it preserves pullbacks (hence \mathbf{C} is required to have pullbacks) and if all naturality squares of the unit and multiplication of the monad are pullbacks.

- Profunctor: A profunctor gives a family of sets of bioperations of fixed arity and coarity.

Arities (or coarities) are governed by the monad T . For example, an operad will be a span / profunctor from $T\mathbf{1}$ to $\mathbf{1}$, where T takes care of the multiplicity of inputs ($\mathbf{1}$ is the category with one object and one morphism). Similarly, a cooperad will go from $\mathbf{1}$ to $T\mathbf{1}$.

The span approach and the profunctor approach should be related, since one goes from profunctors to spans via the “element” or so-called Grothendieck construction. But under this correspondence the (bi)category of profunctors is (bi)equivalent to a subcategory of spans only, the discrete fibrations, while on the other hand the span approach leaves a lot of freedom on the choice of the underlying category \mathbf{C} . As a matter of fact, the two approaches have led to different types of successes. The current state of the art seems to be that:


1. using spans, an impressive variety of shapes in the *non-symmetric case* have been covered. We refer to [[23]] for a book-length account;
2. using profunctors, one may cover the two other kinds of variations mentioned in this introduction.

In the sequel, we use (and introduce) the profunctor road. In Section 2, we introduce several monads on \mathbf{Cat} . After recalling the notion of Kan extension in Section 3, we present two classical categorical accounts of operads in Sections 4 and 5. We proceed then to profunctors in Section 6. A plan of the rest of the paper is given at the end of that section.


Notation We shall use juxtaposition to denote functor application. Moreover, if F is a functor from, say \mathbf{C} to $\mathbf{Set}^{\mathbf{D}}$, we write FXD for $(FX)D$, etc... .

2 Three useful combinators


Consider the following three operations, or combinators. Think of them as morphisms in a category whose objects are (possibly empty) sequences (C_1, \dots, C_n) , where the C_i ’s range over the objects of some category \mathbf{C} .



These combinators should satisfy equations:




(compare with the familiar equations of transpositions, and notice that here we do not have to index them over natural numbers).



The other equations are the familiar ones for comonoids, and (mutatis mutandis) for distributive laws (see below). For a complete list, we refer the reader to [[9, 22]] .


Take any $X \subseteq \{\sigma, \delta, \epsilon\}$. We build a category $!_X \mathbf{C}$ as follows: objects are sequences (C_1, \dots, C_n) of objects of \mathbf{C} .² Morphisms are *string diagrams* built out of the combinators taken from X and out of the morphisms of \mathbf{C} , quotiented over all the equalities that concern the combinators of X , including their naturality, and the equalities



²This notation, that comes from linear logic [[18]] , stresses the idea of multiple input (see [[14]] for explicit links with linear logic).

which allow to embed \mathbf{C} functorially into $!_X \mathbf{C}$. String diagrams are combinations of vertical and horizontal combinations of the basic combinators $\sigma, \delta, \epsilon, f$, such as the ones drawn above.

For six of the eight choices of X , the functor $!_X$ is equipped with a monad structure: the unit takes C to (C) and $f : C \rightarrow D$ to $f : (C) \rightarrow (D)$ and the multiplication on objects is the usual flattening, which takes $((C_1^1, \dots, C_1^{i_1}), (C_n^1, \dots, C_n^{i_n}))$ to $(C_1^1, \dots, C_n^{i_n})$. A little care is needed to define the “flattening” on morphisms. A morphism of $!\mathbf{C}$ is an assembling of boxes connected by combinators typed in $!\mathbf{C}$. When we remove the boxes, we need to turn these combinators into (assemblings of) combinators typed in \mathbf{C} . We call this an expansion. A σ is expanded by means of σ ’s, an ϵ is expanded by putting ϵ ’s in parallel, but we need δ ’s *and* σ ’s to expand a δ . For example, at type (C_1, C_2) , the expansion of $\delta : ((C_1, C_2)) \rightarrow ((C_1, C_2), (C_1, C_2))$ is the following morphism from (C_1, C_2) to (C_1, C_2, C_1, C_2) :



We therefore exclude $!_{\{\delta\}}$ and $!_{\{\delta, \epsilon\}}$ from our treatment. But all the other $!$ constructions are fine.

Three of the six remaining combinations are of particular interest. If $X = \emptyset$ (resp. $X = \{\sigma\}$, $X = \{\sigma, \delta, \epsilon\}$), then $!\mathbf{C}$ is the free strict monoidal (resp. symmetric monoidal, cartesian) category over \mathbf{C} . (By cartesian category, we mean “category with specified finite products”). We write them $!_m, !_s, !_f$, respectively:

monoidal	symmetric monoidal	cartesian
$!_m$	$!_s$	$!_f$

When $X = \{\sigma, \delta, \epsilon\}$, the $!$ -algebras are the cartesian categories, and when $X = \emptyset$ (resp. $X = \{\sigma\}$), the (pseudo-) $!$ -algebras are the monoidal (resp. symmetric monoidal) categories. We refer to [[23]] for details.

We define $?\mathbf{C} = (!(\mathbf{C}^{op}))^{op}$. Graphically, this amounts to reversing the basic combinators σ, δ, ϵ (which could then, if we cared, be called σ, μ, η), while maintaining the orientation of the combinators f imported from \mathbf{C} (since they become themselves again after two op ’s). But it is more convenient to stick with σ, δ, ϵ , to reverse the direction of the f ’s, and to read the diagrams in the bottom-to-top direction. In particular, when $\mathbf{C} = \mathbf{1}$ (the terminal category), there is no f to reverse...


When $X = \emptyset$ or $X = \{\sigma\}$, then $?\mathbf{C}$ is (isomorphic to) $!\mathbf{C}$. When $X = \{\sigma, \delta, \epsilon\}$, $?\mathbf{C}$ is the free cocartesian category over \mathbf{C} . The objects of $?\mathbf{1}$ are

written $0, 1, 2, \dots$, standing for $()$, (\cdot) , (\cdot, \cdot) , \dots , where \cdot is the unique object of $\mathbf{1}$.

Note that for any choice of $?$, there is a faithful functor from $?1$ to **Set**. For any morphism from m to n , i.e., for any string diagram constructed out of the combinators in X that has n input wires and m output wires, one constructs a function from $\{0, \dots, m-1\}$ to $\{0, \dots, n-1\}$, by the following rules.

- σ : the transposition on $\{0, 1\}$;
- ϵ : the unique function from the empty set to $\{0\}$;
- δ : the unique function from $\{0, 1\}$ to $\{0\}$;
- vertical composition of string diagrams: function composition;
- horizontal composition of string diagrams: their categorical sum, i.e., for $f : \{0, \dots, m-1\} \rightarrow \{0, \dots, n-1\}$ and $g : \{0, \dots, p-1\} \rightarrow \{0, \dots, q-1\}$, $f+g : \{0, \dots, m+p-1\} \rightarrow \{0, \dots, n+q-1\}$ is defined by $(f+g)(i) = f(i)$ if $i < m$ and $f+g(i) = g(i) + n$ if $i \geq m$.

More synthetically, the function f associated with a string diagram is obtained by naming the output and input wires $0, 1, \dots, m-1$ and $0, 1, \dots, n-1$, respectively, and then computing $f(i)$ as the name of the input wire reached when starting from input wire i , going up in δ nodes, and going up left or right in σ nodes according to whether this node is reached from down right or down left, respectively. For example, the picture



represents the function $f : \{0, 1, 2\} \rightarrow \{0, 1, 2\}$ defined by

$$f(0) = f(1) = 1 \quad f(2) = 0$$

Note on this example that ϵ witnesses the lack of surjectivity. More precisely, as first shown by Burroni [[9]] , the functor, when suitably corestricted, gives the following isomorphisms and equivalences of categories:

$?_{\emptyset}$	identity functions
$?_{\{\sigma\}}$	bijections
$?_{\{\sigma, \delta, \epsilon\}}$	all functions
$?_{\{\delta, \epsilon\}}$	monotone functions
$?_{\{\sigma, \delta\}}$	surjective functions
$?_{\{\sigma, \epsilon\}}$	injective functions

where, say, the last line should be read as follows: the category $?_{\{\sigma, \epsilon\}}(\mathbf{1})$ is equivalent to the category of finite sets and injective functions, and isomorphic to its full subcategory whose objects are the sets $\{0, \dots, i-1\}$ ($i \in \mathbb{N}$). For $?_{\{\delta, \epsilon\}}(\mathbf{1})$, one must take the category of all finite total orders.

3 Kan extensions

We recall that given a functor $K : \mathbf{M} \rightarrow \mathbf{C}$, a left Kan extension of a functor $T : \mathbf{M} \rightarrow \mathbf{A}$ along K is a pair of a functor $Lan_K(T) : \mathbf{C} \rightarrow \mathbf{A}$ and a natural transformation $\eta_T : T \rightarrow Lan_K(T)K$ such that $(Lan_K(T), \eta_T)$ is universal from T to \mathbf{A}^K . This means that for any other pair of a functor $S : \mathbf{C} \rightarrow \mathbf{A}$ and a natural transformation $\alpha : T \rightarrow SK$ there exists a unique natural transformation $\mu : Lan_K(T) \rightarrow S$ such that $\alpha = \mu K \circ \eta$. Here we shall only need to know that the solution of this universal problem is given by the following formula for $Lan_K(T)$:

$$Lan_K TC = \int^m \mathbf{C}[KM, C] \cdot TM$$

where we use Mac Lane's notation for coends [[24]]. Coends are sorts of colimits (or inductive limits), adapted to the case of diagrams which vary both covariantly and contravariantly over some parameter: here, M appears contravariantly in $\mathbf{C}[KM, C]$ and covariantly in TM . In the formula, $\mathbf{C}[KM, C] \cdot TM$ stands for the coproduct of as many copies of TM as there are morphisms from KM to C .

$$\begin{array}{ccc} \mathbf{M} & \xrightarrow{K} & \mathbf{C} \\ \downarrow T & \nearrow Lan_K T & \\ \mathbf{A} & & \end{array}$$

When K is full and faithful, then η is iso, and in particular the triangle commutes (up to isomorphism). When moreover K is the Yoneda embedding, we get

$$Lan_Y TX = \int^M Xm \cdot TM$$

When moreover \mathbf{A} is a presheaf category $\mathbf{Set}^{\mathbf{D}}$, we have (since limits and colimits of presheaves are pointwise):

$$Lan_Y TXD = \int^M XM \times TMD$$

or, making the quotient involved in the coend explicit:

$$Lan_Y TXD = (\sum_M XM \times TMD) / \approx$$

When \approx is termwise, i.e. is the disjoint union of equivalences \approx_M each on $XM \times TMD$, we get a formula which will look more familiar to algebraic operadists:

$$Lan_{\mathcal{Y}} T X D = \sum_M (XM \otimes_M TMD)$$

(where we have written $(XM \times TMD)/\approx_M$ as $XM \otimes_M TMD$). In our examples, \approx will not always be termwise (it will be in the operad cases, but not in the clone case).

4 Kelly's account of operads

In 1972, Kelly gave the following description of operads [[21]] (see also [[13]]):

$$\begin{array}{ccccc} \mathbf{1} & \xrightarrow{\quad} & !\mathbf{1} & \xrightarrow{\quad \mathcal{Y} \quad} & \mathbf{Set}^{?\mathbf{1}} \\ & \searrow X & \downarrow X^{\otimes -} & \swarrow Lan_{\mathcal{Y}}(X^{\otimes -}) = \bullet X & \\ & & \mathbf{Set}^{?\mathbf{1}} & & \end{array}$$

In this diagram, the functor $X^{\otimes -}$ associates with n the n th iterated tensor of X , with respect to the following tensor product structure on $\mathbf{Set}^{?\mathbf{1}}$ due (in a more general setting) to Day [[12]]:

$$(X \otimes Y)p = \int^{m,n} X m \times Y n \times ?\mathbf{1}[m+n, p]$$

The n -ary tensor product is described by the following formula (which also gives the unit of the tensor product, for $m = 0$):

$$(X_1 \otimes X_2 \otimes \dots \otimes X_m)p = \int^{n_1, \dots, n_m} X_1 n_1 \times \dots \times X_m n_m \times ?\mathbf{1}[n_1 + \dots + n_m, p]$$

Hence the formula for $Lan_{\mathcal{Y}}(X^{\otimes -})Y$, which we write $Y \bullet X$, is

$$\begin{aligned} (Y \bullet X)p &= \int^m Y m \times X^{\otimes m} p \\ &= \int^{m, n_1, \dots, n_m} Y m \times X n_1 \times \dots \times X n_m \times ?\mathbf{1}[n_1 + \dots + n_m, p] \end{aligned}$$

In the case of $!_f$ this boils down to

$$(Y \bullet X)p = \int^m Y m \times (Xp)^m$$

This is because in a cocartesian category, a morphism $f : n_1 + \dots + n_m \rightarrow p$ amounts to morphisms $f_i \in ?\mathbf{1}[n_i, p]$ which allows us to define a map from $X n_1 \times \dots \times X n_m \times ?\mathbf{1}[n_1 + \dots + n_m, p]$ to $(Xp)^m$ as follows:

$$(a_1, \dots, a_m, f) \mapsto (X f_1 a_1, \dots, X f_m a_m)$$

and then to “extend” cocones indexed over m to cocones indexed over m, n_1, \dots, n_m .

One can check rather easily that the *substitutions operation* \bullet , together with the unit defined by

$$In = ?\mathbf{1}[1, n]$$

provide a (non-symmetric) monoidal structure on $\mathbf{Set}^{?1}$ (for $I \bullet X \approx X$, one uses the fact that X^{\otimes} - is functorial).

When $!$ is $!_m, !_s, !_f$, respectively, a monoid for this structure (I, \bullet) is an operad, a symmetric operad, a clone, respectively.

5 Operads from analytic functors

Another way to arrive at the operation \bullet just defined is via a different Kan extension:

$$\begin{array}{ccc} ?\mathbf{1} & \xrightarrow{\subseteq} & \mathbf{Set} \\ \downarrow X & \nearrow Lan_{\subseteq} X & \\ \mathbf{Set} & & \end{array}$$

where \subseteq is the faithful (and non full) functor described at the end of Section 2. This is the approach taken by Joyal (for $!_s$) [[20]] . In Joyal's language, X is called a *species of structure*, and $Lan_{\subseteq} X$ is the associated *analytic functor*, whose explicit formula is (for any set z):

$$Lan_{\subseteq} Xz = \int^m z^m \times Xm$$

It can be shown that Lan_{\subseteq} is faithful. Then $Y \bullet X$ is characterised by the following equality:

$$Lan_{\subseteq}(Y \bullet X) = Lan_{\subseteq} Y \circ Lan_{\subseteq} X$$

which evidences the fact that \bullet is a composition operation. Indeed, we have:

$$\begin{aligned} & Lan_{\subseteq} Y(Lan_{\subseteq} Xz) \\ &= \int^m (\int^n z^n \times Xn)^m \times Ym \\ &= \int^{m, n_1, \dots, n_m} z^{n_1 + \dots + n_m} \times Xn_1 \times \dots \times Xn_m \times Ym \\ &= \int^p z^p \times (\int^{m, n_1, \dots, n_m} Ym \times Xn_1 \times \dots \times Xn_m \times ?\mathbf{1}[n_1 + \dots + n_m, p]) \\ &= Lan_{\subseteq}(Y \bullet X)z \end{aligned}$$

(the summation over p is superfluous since from $f \in ?\mathbf{1}[n_1 + \dots + n_m, p]$ we get $z^{\subseteq(f)}$ from z^p to $z^{n_1 + \dots + n_m}$).

6 Profunctors

Recall that a profunctor (or distributor) [[5, 6]] Φ from \mathbf{C} to \mathbf{C}' is a functor

$$\Phi : \mathbf{C} \times \mathbf{C}'^{op} \rightarrow \mathbf{Set}$$

We write $\Phi : \mathbf{C} \multimap \mathbf{C}'$. Composition of profunctors is given by the following formula:

$$(\Psi \circ \Phi)(C, C'') = \int^{C'} \Psi(C', C'') \times \Phi(C, C')$$

Therefore, profunctors compose only up to isomorphism. Categories, profunctors, and natural transformations form thus, not a 2-category, but a bicategory [[4]] .

The bicategory **Prof** of profunctors is self-dual, via the isomorphism $^{op} : \mathbf{Prof} \rightarrow \mathbf{Prof}^{op}$ which maps \mathbf{C} to \mathbf{C}^{op} and $\Phi : \mathbf{C}_1 \multimap \mathbf{C}_2$ to

$$\Phi^{op} = ((C_2, C_1) \mapsto \Phi(C_1, C_2)) : \mathbf{C}_2^{op} \multimap \mathbf{C}_1^{op}$$

The composition of profunctors can be synthesised via Kan extensions: Given $\Phi : \mathbf{C} \multimap \mathbf{C}'$ and $\Psi : \mathbf{C}' \multimap \mathbf{C}''$, consider their “twisted curried”³ versions $\Phi' : \mathbf{C}'^{op} \rightarrow \mathbf{Set}^{\mathbf{C}}$ and $\Psi' : \mathbf{C}''^{op} \rightarrow \mathbf{Set}^{\mathbf{C}'}$, defined by $\Phi' C' C = \Phi(C, C')$ and $\Psi' C'' C' = \Psi(C', C'')$. Then it is immediate to check that $\Psi \circ \Phi$ is the uncurried version of $(Lan_{\mathcal{Y}} \Phi') \circ \Psi'$, as illustrated in the following diagram:

$$\begin{array}{ccc} \mathbf{C}'^{op} & \xrightarrow{\mathcal{Y}} & \mathbf{Set}^{\mathbf{C}'} \xleftarrow{\Psi'} \mathbf{C}''^{op} \\ \downarrow \Phi' & \swarrow Lan_{\mathcal{Y}} \Phi' & \searrow (\Psi \circ \Phi)' \\ \mathbf{Set}^{\mathbf{C}} & & \end{array}$$

Now we take “profunctor glasses” to look at Kelly’s and Joyal’s diagrams.

- A presheaf $X : ?\mathbf{1} \rightarrow \mathbf{Set}$ can be viewed as a profunctor

$$(n \mapsto (\cdot \mapsto Xn)) : ?\mathbf{1} \multimap \mathbf{1}$$

- $X^{\otimes -}$ can be viewed as a profunctor from $?\mathbf{1}$ to $?\mathbf{1}$ (since $(?\mathbf{1})^{op} = !(\mathbf{1}^{op}) = !\mathbf{1}$). Thus the transformation

$$\frac{X : ?\mathbf{1} \multimap \mathbf{1}}{X^{\otimes -} : ?\mathbf{1} \multimap ?\mathbf{1}}$$

suggests to consider $?$ as a comonad over **Prof**.

- The diagram and the formula defining $Y \bullet X$ in Section 4 exhibit $Y \bullet X$ as the profunctor composition of $Y : ?\mathbf{1} \multimap \mathbf{1}$ and $X^{\otimes -} : ?\mathbf{1} \multimap ?\mathbf{1}$. But a better way to put it is that $Y \bullet X$ is the composition of Y and X in the coKleisli bicategory **Prof**_?, and we shall see that it is indeed the case.

³After the name of Curry, who defined a calculus of functions called combinatory logic, based on application and a few combinators, and where functions of several arguments are expressed through repeated applications.

- Joyal’s construction amounts to “taking points”. We first observe that $? \emptyset = \mathbf{1}$, where \emptyset is the empty category. (This holds for any of the eight choices for $!$.) It follows that \emptyset is terminal in $\mathbf{Prof}_?$. In this category, the points of an object \mathbf{C} are the morphisms from \emptyset to \mathbf{C} , i.e., the profunctors from $\mathbf{1}$ to \mathbf{C} , i.e. the presheaves over \mathbf{C} . In particular, the points of $\mathbf{1}$ are just sets. We shall see that

$$\mathbf{Prof}_?[\emptyset, X] = \text{Lan}_{\subseteq} X$$

i.e., that the analytic functor associated with the presheaf X describes its pointwise behaviour in $\mathbf{Prof}_?$.

Our goal is thus to figure out $?$ as a comonad on \mathbf{Prof} . We shall do this in three steps:

- In Section 7, we show that profunctors arise as a Kleisli category for the presheaf construction $\mathbf{C} \mapsto \mathbf{Set}^{\mathbf{C}^{op}}$.
- In Section 9, we show that all the monads $!$ of Section 2 distribute over Psh . Distributive laws are recalled in Section 8. In Section 10, we pause to explain how both Day’s tensor product and this distributive law can be synthesised out of considerations of structure preservation.
- in Section 11, we show that this distributive law allows us to extend $!$ to a monad on \mathbf{Prof} , and by self-duality we obtain the comonad $?$ that we are looking for.

Warning. In what follows, we ignore coherence issues for simplicity, and we partially address size issues.

- Coherence issues arise from the fact that we shall compose morphisms using coend formulas, which make sense only up to iso. In particular, *our “distributive law” will in fact be a “pseudo-distributive law”* [[25, 10, 16]] .
- Size issues arise from the fact that the presheaf construction is dramatically size increasing. It is therefore in fact simply not rigorous to call Psh a monad (or even a pseudo-monad) on \mathbf{Cat} . But the issue is fortunately not too severe. In a forthcoming paper, Fiore, Gambino, Hyland, and Winskel propose a general notion of Kleisli structure, which we shall sketch here (ignoring again coherence issues), and in which Psh fits.

7 Profunctors as a Kleisli category

If we write a profunctor $\Phi : \mathbf{C}_1 \multimap \mathbf{C}_2$ in (untwisted) curried form

$$C_1 \mapsto (C_2 \mapsto \Phi(C_1, C_2)) : \mathbf{C}_1 \rightarrow \mathbf{Set}^{\mathbf{C}_2^{op}}$$

this suggests us to look at the operation Psh on the objects of \mathbf{Cat} defined by $Psh(\mathbf{C}) = \mathbf{Set}^{\mathbf{C}^{op}}$. The idea is to exhibit Psh as a (pseudo-)monad, so that profunctors arise as a Kleisli category. We have a good candidate for the unit η , namely: $\mathcal{Y} : \mathbf{C} \rightarrow Psh(\mathbf{C})$.

But the Yoneda functor makes sense only for a locally small category (one in which all homsets are sets), while it is not clear at all whether Psh keeps us within the realm of locally small categories, i.e. if \mathbf{C} is locally small, then $\mathbf{Set}^{\mathbf{C}^{op}}$ is not necessarily locally small. It is nevertheless tempting to go on with a multiplication $\mu : \mathbf{Set}^{(\mathbf{Set}^{\mathbf{C}^{op}})^{op}} \rightarrow \mathbf{Set}^{\mathbf{C}^{op}}$ given by a left Kan extension

$$\begin{array}{ccc} \mathbf{Set}^{\mathbf{C}^{op}} & \xrightarrow{\mathcal{Y}} & \mathbf{Set}^{(\mathbf{Set}^{\mathbf{C}^{op}})^{op}} \\ id \downarrow & \swarrow \mu = Lan_{\mathcal{Y}}(id) & \\ \mathbf{Set}^{\mathbf{C}^{op}} & & \end{array}$$

with explicit formula $\mu(H)C = \int^F HF \times FC$. But why should a coend on such a vertiginous indexing collection exist?

We pause here to recall that an equivalent presentation of a monad on a category \mathbf{C} is by means of the following data:

- for each object C of \mathbf{C} , an object TC of \mathbf{C} , and a morphism $\eta_C \in \mathbf{C}[C, TC]$;
- for all objects C, D of \mathbf{C} and each morphism $f \in \mathbf{C}[C, TD]$, a morphism $f^\# \in \mathbf{C}[TC, TD]$;

satisfying the equations $f = f^\# \circ \eta = f, \eta^\# = id, (g^\# \circ f)^\# = g^\# \circ f^\#$.

Let us also recall the definition of the associated Kleisli category \mathbf{C}_T : its objects are the objects of \mathbf{C} , and one sets $\mathbf{C}_T[C, D] = \mathbf{C}[C, TD]$, with composition easily defined using the composition in \mathbf{C} and the $_^\#$ operation.

It turns out that under this guise, the definition of monad can be generalised in a way that will fit our purposes. Ignoring coherence issues, a *Kleisli structure* (as proposed in [[15]]) is given by the following data:

- a collection \mathcal{A} of objects of \mathbf{C} ;
- for each object A in \mathcal{A} an object TA of \mathbf{C} , and a morphism $\eta_A \in \mathbf{C}[A, TA]$;
- for all objects A, B in \mathcal{A} and each morphism $f \in \mathbf{C}[A, TB]$, a morphism $f^\# \in \mathbf{C}[TA, TB]$;

satisfying the same equations. The associated Kleisli category \mathbf{C}_T has \mathcal{A} as collection of objects, and one sets $\mathbf{C}_T[A, B] = \mathbf{C}[A, TB]$. We recover the monads when every object of \mathbf{C} is in \mathcal{A} . But in general there is no such thing as a multiplication, since it is not granted that TTC exists for all C . We also note that in a Kleisli structure, T is still a functor, not from \mathbf{C} to \mathbf{C} , but from $\mathbf{C} \downarrow_{\mathcal{A}}$ (the full subcategory spanned by \mathcal{A}) to \mathbf{C} .

In our setting, we have $\mathbf{C} = \mathbf{Cat}$, and we can take \mathcal{A} to consist of the small categories (in which the objects form a set, as well as all homsets). Then we have no worry about the coends that we shall write. We complete the definition of the Kleisli structure with the definition of the $\#$ operation.

$$\begin{array}{ccc} \mathbf{C} & \xrightarrow{y} & \mathbf{Set}^{\mathbf{C}^{op}} \\ F \downarrow & \swarrow & \\ \mathbf{Set}^{\mathbf{D}^{op}} & & F^\# = \text{Lan}_y(F) \end{array}$$

and the explicit formula is $F^\#XD = \int^C XC \times FCD$. It is then an easy exercise to check that the three equations of a Kleisli structure are satisfied, and that the composition in the associated Kleisli category coincides with the composition of profunctors as defined in the previous section.

We end the section with a description of the functorial action of Psh (on, say, functors between small categories):

$$Psh(F)XD = (\eta_{Psh} \circ F)^{\#}XD = \int^C XC \times \mathbf{D}[D, FC]$$


8 Distributive laws

Recall that a distributive law $[[3]]$ (see also $[[2]]$) is a natural transformation $\lambda : TS \rightarrow ST$, where (S, η_S, μ_S) and (T, η_T, μ_T) are two monads over the same category \mathbf{C} , satisfying the following laws, expressed in the language of string diagrams:

EQUATION $(\lambda - \eta_T)$:


The diagram shows an equality between two expressions. On the left, a node labeled λ has four incident lines: two labeled T (top-left and bottom-right) and two labeled S (top-right and bottom-left). A line labeled S enters the top-left T line from above. A line labeled T enters the top-right S line from above. This structure is connected to an oval node labeled η . On the right, there is a single vertical line labeled T at the bottom, connected to an oval node labeled η . To the right of this is a vertical line labeled S .

EQUATION $(\lambda - \mu_T)$:



and two similar equations $(\lambda - \mu_S)$ and $(\lambda - \eta_S)$.

A distributive law allows us to extend T to \mathbf{C}_S , the Kleisli category of S (and conversely, such a lifting induces a distributive law such that the two constructions are inverse to each other).⁴ The extended T acts on objects as the old T . On morphisms, its action is described as follows:



where the box separates the inside which lives in \mathbf{C} from the outside which lives in \mathbf{C}_S . It is a nice exercise to check that the old and the new T satisfy

$$T \circ F_S = F_S \circ T$$

where $F_S : \mathbf{C} \rightarrow \mathbf{C}_S$ is defined by $F_S(C) = C$ and $F_S(f) = \eta_S \circ f$, and that the new T is a monad.

⁴ From another standpoint, a distributive law $\lambda : TS \rightarrow ST$ also amounts to lifting S to the category of T -algebras. See Section 10.

In the next section, we want to apply this to $S = Psh$ and $T = !$. But S is only a Kleisli structure. Only slight adjustments is needed:

1. make sure that \mathcal{A} is stable under T ;
2. replace the equation $(\lambda - \mu_S)$ by the following one (for $f : A \rightarrow SB$):

$$\lambda_B \circ T(g^\#) = (\lambda_B \circ Tg)^\# \circ \lambda_A$$

9 $A \text{ !}/Psh$ distributive law

We define a transformation $\lambda : ! \circ Psh \rightarrow Psh \circ !$, as follows:

- on objects:

$$(F_1, \dots, F_n) \mapsto ((C_1, \dots, C_m) \mapsto \int^{D_1, \dots, D_n} F_1 D_1 \times \dots \times F_n D_n \times !\mathbf{C}[(C_1, \dots, C_m), (D_1, \dots, D_n)])$$

- on generating morphisms:

- $\alpha : F \rightarrow G$. There is an obvious map from $\int^D FD \times !\mathbf{C}[C_1, \dots, C_m, (D)]$ to $\int^D GD \times !\mathbf{C}[C_1, \dots, C_m, (D)]$.
- Combinators, say $\delta : (F) \rightarrow (F, F)$. We obtain a map

$$\begin{array}{l} \text{from } \int^D FD \times !\mathbf{C}[C_1, \dots, C_m, (D)] \\ \text{to } \int^{D_1, D_2} FD_1 \times FD_2 \times !\mathbf{C}[(C_1, \dots, C_m), (D_1, D_2)] \end{array}$$

by going

$$\begin{array}{l} \text{from } FD \times !\mathbf{C}[C_1, \dots, C_m, (D)] \\ \text{to } FD \times FD \times !\mathbf{C}[(C_1, \dots, C_m), (D, D)] \end{array}$$

We verify three of the four equations (leaving the last one – as stated at the end of Section 8 – to the reader):

$(\lambda - \eta_l)$. The left hand side is computed by taking the case $n = 1$ in the definition of λ , thus we obtain

$$F \mapsto ((C_1, \dots, C_m) \mapsto \int^D FD \times !\mathbf{C}[(C_1, \dots, C_m), (D)])$$

which is the formula for $Psh(\eta_l)$.

$(\lambda - \eta_{Psh})$. We have (at $(A_1, \dots, A_n), (C_1, \dots, C_m)$):

$$\begin{aligned} & \int^{D_1, \dots, D_n} \mathbf{C}[D_1, A_1] \times \dots \times \mathbf{C}[D_n, A_n] \times !\mathbf{C}[(C_1, \dots, C_m), (D_1, \dots, D_n)] \\ & = !\mathbf{C}[(C_1, \dots, C_m), (A_1, \dots, A_n)] = \mathcal{Y}(A_1, \dots, A_n)(C_1, \dots, C_m) \end{aligned}$$

$(\lambda - \mu_!)$. The left hand side is (at $((F_1^1, \dots, F_1^{i_1}), \dots, (F_n^1, \dots, F_n^{i_n})), (C_1, \dots, C_m))$):

$$\int^{D_1^1, \dots, D_n^{i_n}} F_1^1 D_1^1 \times \dots \times F_n^{i_n} D_n^{i_n} \times !\mathbf{C}[\vec{C}, (D_1^1, \dots, D_n^{i_n})]$$

We first compute the upper part $(\lambda!) \circ (!\lambda)$ of the right hand side (at $(\vec{F}_1, \dots, \vec{F}_n), ((A_1^1, \dots, A_1^{j_1}), \dots, (A_p^1, \dots, A_p^{j_p}))$):

$$\begin{aligned} & \int^{\vec{B}_1, \dots, \vec{B}_n} \int^{D_1^1, \dots, D_n^{i_n}} \vec{F}_1^1 D_1^1 \times \dots \times F_n^{i_n} D_n^{i_n} \times !\mathbf{C}[\vec{B}_1, \vec{D}_1] \times \dots \times !\mathbf{C}[\vec{C}_n, \vec{D}_n] \times !!\mathbf{C}[\vec{A}, \vec{B}] \\ &= \int^{D_1^1, \dots, D_n^{i_n}} F_1^1 D_1^1 \times \dots \times F_n^{i_n} D_n^{i_n} \times !!\mathbf{C}[\vec{A}, \vec{D}] \end{aligned}$$

Finally, applying $Psh(\mu)$, we get (at \vec{F}, \vec{C}):

$$\int^{\vec{A}_1, \dots, \vec{A}_p} \int^{D_1^1, \dots, D_n^{i_n}} F_1^1 D_1^1 \times \dots \times F_n^{i_n} D_n^{i_n} \times !!\mathbf{C}[\vec{A}, \vec{D}] \times !\mathbf{C}[\vec{C}, (A_1^1, \dots, A_p^{j_p})]$$

and we conclude using $\mu_!$.

Summarizing, we have the following result:

Proposition 9.1 *The transformation λ defines a (pseudo)-distributive law for any choice among the six $!_X$ monads over the presheaf Kleisli structure.*

A similar proposition is proved in [[26]] : there, the existence of λ is derived from the fact $!$ and Psh are both free constructions (the latter being the free cocompletion, i.e. the free category with all colimits) and commute in the sense that Psh lifts to the category of $!$ -algebras (cf. Section 2). For completeness, we sketch this more conceptual approach in the following section.

On the other hand, the advantage of a “symbol-pushing” proof like the one presented above is to highlight plainly the “uniformity” in the choice of any combination X of Burroni’s combinators.

10 Intermezzo

The reader may have noticed that the formula defining λ proposed in the last section “looks like” the formula for Day’s tensor product (cf. Section 4). In this section, we actually synthesise the latter formula from considerations of cocontinuity, and then the former one from considerations of monad lifting. Since this section has merely an explanatory purpose, we limit ourselves here to the case $! = !_\emptyset$, and we disregard size issues.

Cocontinuity is at the heart of the presheaf construction. It is well-known that $\mathbf{Set}^{\mathbf{C}^{op}}$ is the free cocomplete category over \mathbf{C} , and that the unique cocontinuous (i.e., colimit preserving) extension of a functor $F : \mathbf{C} \rightarrow \mathbf{D}$ (where \mathbf{D} is cocomplete, i.e., has all colimits) is its left Kan extension $Lan_{\mathbf{Y}}(F)$:

$$\begin{array}{ccc}
\mathbf{C} & \xrightarrow{\mathcal{Y}} & \mathbf{Set}^{\mathbf{C}^{op}} \\
\downarrow F & \swarrow Lan_{\mathcal{Y}} F & \\
\mathbf{D} & &
\end{array}$$

In particular, the fact that $Lan_{\mathcal{Y}}(F)$ preserves colimits follows from its being left adjoint to $(C \mapsto (D \mapsto \mathbf{D}[FC, D]))$ (easy check).

We could have defined the “monad” Psh more economically and more conceptually as the monad arising from the adjunction between \mathbf{Cat} and the category of cocomplete categories. The “brute force” construction of last section is in any case a good exercise in coend computations.

Although we shall not need it here, it is worth pointing out that the adjunction is monadic, i.e., that the (pseudo-) Psh -algebras are actually the cocomplete categories (and hence that there can be at most one Psh -algebra structure on a given category). This can be proved using Beck’s characterization of monadic adjunctions (Theorem 4.4.4 of [[7]]).

Consider a slightly more general version of Day’s tensor product than that given in Section 4, with now some monoidal category \mathbf{C} in place of $\mathbf{1}$

$$(X \otimes Y)C = \int^{C_1, C_2} XC_1 \times XC_2 \times \mathbf{C}[C, C_1 \otimes C_2]$$

(thus Day’s tensor depends on the tensor on \mathbf{C}). We show that this definition is entirely determined from the requirements that

- \otimes extends the tensor product of \mathbf{C} (via the Yoneda embedding), and
- \otimes is cocontinuous in each argument.

And, of course, Day’s product does satisfy these requirements (easy proof left to the reader). Recall that \mathcal{Y} is dense, i.e., that every presheaf is a colimit of representable presheaves (i.e. presheaves of the form YC), and, more precisely, that $Lan_{\mathcal{Y}}\mathcal{Y} = id$:

$$X = \int^D XD \cdot \mathcal{Y}D = colim_{(D, z \in XD)} \mathcal{Y}D$$

Then, we must have

$$\begin{aligned}
X \otimes Y &= (colim_{(D_1, z \in XD_1)} \mathcal{Y}D_1) \otimes Y \\
&= colim_{(D_1, z \in XD_1)} (\mathcal{Y}D_1 \otimes Y) && \text{(by requirement 2)} \\
&= colim_{(D_1, z \in XD_1), (D_2, z \in XD_2)} (\mathcal{Y}D_1 \otimes \mathcal{Y}D_2) && \text{(by requirement 2)} \\
&= colim_{(D_1, z \in XD_1), (D_2, z \in XD_2)} \mathcal{Y}(D_1 \otimes D_2) && \text{(by requirement 1)} \\
&= \int^{D_1, D_2} (XD_1 \times XD_2) \cdot \mathcal{Y}(D_1 \otimes D_2)
\end{aligned}$$

In other words, Day’s tensor is determined, as announced.

It is shown in [[19]] that the (2-) adjunction between categories and cocomplete categories specialises to an adjunction between monoidal categories and

cocomplete monoidal categories, and that $\mathbf{Set}^{C^{op}}$, equipped with Day's tensor product, is the free such one. We content ourselves here with the key verification, namely that when $F : \mathbf{C} \rightarrow \mathbf{D}$ is monoidal, then so is its unique cocontinuous extension $\tilde{F} = Lan_{\mathcal{Y}} F$. We first compute $\tilde{F}X_1 \otimes \tilde{F}X_2$ and $\tilde{F}(X_1 \otimes X_2)$:

$$\begin{aligned}\tilde{F}X_1 \otimes \tilde{F}X_2 &= \int^{C_1, C_2} (X_1 C_1 \times X_2 C_2) \cdot (FC_1 \otimes FC_2) \quad (\text{by cocontinuity}) \\ \tilde{F}(X_1 \otimes X_2) &= \int^C \int^{C_1, C_2} X_1 C_1 \times X_2 C_2 \times \mathbf{C}[C, C_1 \otimes C_2] \cdot FC \\ &= \int^{C_1, C_2} X_1 C_1 \times X_2 C_2 \cdot F(C_1 \otimes C_2)\end{aligned}$$

From this, it follows that the maps $FC_1 \otimes FC_2 \rightarrow F(C_1 \otimes C_2)$ induce a map $\tilde{F}X_1 \otimes \tilde{F}X_2 \rightarrow \tilde{F}(X_1 \otimes X_2)$.

As a consequence Psh lifts to (pseudo-) $!$ -algebras, i.e. to monoidal categories. Given two monads T, S , a lifting of S to the category of T -algebras consists of a monad S' on the category of T -algebras such that S, S' commute with the forgetful functor, and such that the unit and the multiplication of S' are mapped by the forgetful functor to the unit and the multiplication of S . Here, we take $S = Psh$ and $T = !$. The lifting S' maps a monoidal structure \otimes on \mathbf{C} to the corresponding Day's monoidal structure on $Psh(\mathbf{C})$.

We turn back to the general situation. A lifting of S to T -algebras is equivalent to giving a distributive law $\lambda : TS \rightarrow ST$ (cf. Footnote 4). One defines λ from the data of the lifting as follows:

$$\lambda_C = S'(\mu_C) \circ TS\eta_C \quad (\mu, \eta \text{ relative to } T)$$

(recall that μ_C is a T -algebra – the free one). This allows us to derive the distributive law from Day's tensor product. First, for $S = Psh$ and $T = !$, $S'(\mu_C)$ is the Day's tensor product $(G_1, \dots, G_n) \mapsto G_1 \otimes \dots \otimes G_n$ associated with the (flattening) tensor product on $!\mathbf{C}$:

$$(G_1 \otimes \dots \otimes G_n) \vec{C} = \int^{\vec{A}_1, \dots, \vec{A}_n} G_1 \vec{A}_1 \times \dots \times G_n \vec{A}_n \times !\mathbf{C}[\vec{C}, (A_1^1, \dots, A_n^{i_n})]$$

Then λ is obtained by replacing G_i by $Psh(\eta)(F_i)$:

$$\int^{\vec{A}_1, \dots, \vec{A}_n} z_1 \times \dots \times z_n \times !\mathbf{C}[\vec{C}, (A_1^1, \dots, A_n^{i_n})]$$

with $z_1 = \int^{D_1} !F_1 D_1 \times !\mathbf{C}[\vec{A}_1, (D_1)]$, \dots , $z_n = \int^{D_n} F_n D_n \times !\mathbf{C}[\vec{A}_n, (D_n)]$. This simplifies to the formula given in Section 9.

11 The (bi)category \mathbf{Prof} ?

We next consider the coKleisli bicategory $\mathbf{Prof}_? = (\mathbf{Cat}_{Psh})_?$. From the previous sections we know that the monad $!$ on \mathbf{Cat} extends to a (pseudo-) monad $!$ on \mathbf{Prof} , which under the self-duality of \mathbf{Prof} gives a (pseudo-) comonad $?$

on **Prof**. Just as a monad gives rise to a Kleisli category, a comonad gives rise to a coKleisli category. Here, **Prof**_? has categories as objects, while its (1-)morphisms are defined by

$$\mathbf{Prof}_?[\mathbf{C}, \mathbf{C}'] = \mathbf{Prof}[? \mathbf{C}, \mathbf{C}']$$

For $X = \{\sigma\}$, this is the category generalised species of structures of Fiore, Gambino, Hyland, and Winskel [[14]]. Also with $X = \{\sigma\}$, the endomorphisms in **Prof**_? are the C -profiles of Baez and Dolan [[1]].⁵ We are now

in a position to round the circle and to show (cf. Theorem 4.1 of [[14]] for the case $X = \{\sigma\}$) that the composition in this category coincides with the substitution operation as defined in Section 4. We decompose all the steps of the construction.

(1) Let $\Phi : \mathbf{C} \multimap \mathbf{C}'$. Then $! \Phi : ! \mathbf{C} \multimap ! \mathbf{C}'$ is (using λ):

$$! \Phi((C_1, \dots, C_n), \vec{C}') = \int^{D'_1, \dots, D'_n} \Phi C_1 D'_1 \times \dots \times \Phi C_n D'_n \times ! \mathbf{C}'[\vec{C}', \vec{D}']$$

(1)^{op} We can then define $? \Phi = (! \Phi^{op})^{op}$:

$$? \Phi(\vec{C}, (C'_1, \dots, C'_p)) = \int^{D_1, \dots, D_p} \Phi D_1 C'_1 \times \dots \times \Phi D_p C'_p \times ? \mathbf{C}[\vec{D}, \vec{C}]$$

(2) The multiplication of $!$ viewed as a monad on **Prof** at \mathbf{C} is

$$(((C_1^1, \dots, C_1^{i_1}), \dots, (C_n^1, \dots, C_n^{i_n})), \vec{D}) \mapsto ! \mathbf{C}[\vec{D}, (C_1^1, \dots, C_n^{i_n})] : ! \mathbf{C} \multimap ! \mathbf{C}$$

(2)^{op} Hence the comultiplication $\delta : ? \mathbf{C} \multimap ?? \mathbf{C}$ of $? = (\mathbf{C} \mapsto (! \mathbf{C}^{op})^{op})$ is

$$((\vec{C}, ((D_1^1, \dots, D_1^{i_1}), \dots, (D_n^1, \dots, D_n^{i_n}))) \mapsto ? \mathbf{C}[(D_1^1, \dots, D_n^{i_n}), \vec{C}])$$

(3) Let $\Phi : ? \mathbf{C} \multimap \mathbf{C}'$. Then $\Phi^b = (? \Phi) \circ \delta : ? \mathbf{C} \multimap ? \mathbf{C}'$ is given by the following explicit formula (on $\vec{C}, (C'_1, \dots, C'_p)$):

$$\begin{aligned} & \int^{\vec{A}_1, \dots, \vec{A}_m} ? \mathbf{C}[(A_1^1, \dots, A_m^{j_m}), \vec{C}] \times \left(\int^{\vec{D}_1, \dots, \vec{D}_p} \Phi \vec{D}_1 C'_1 \times \dots \times \Phi \vec{D}_p C'_p \times ?? \mathbf{C}[\vec{D}, \vec{A}] \right) \\ &= \int^{\vec{D}_1, \dots, \vec{D}_p} \Phi \vec{D}_1 C'_1 \times \dots \times \Phi \vec{D}_p C'_p \times ? \mathbf{C}[(D_1^1, \dots, D_p^{i_p}), \vec{C}] \end{aligned}$$

⁵ Baez and Dolan use a variation to synthesise the composition of \mathbf{C} -signatures: they note that **Set**^{?C} is a free symmetric monoidal cocomplete construction (where the tensor preserves the colimits in each argument, cf. Section 10), which allows them to identify C -profiles with the endofunctors on **Set**^{?C} that preserve tensor and colimits, and hence to inherit composition from usual composition of functors.

The simplification comes from the functoriality of the multiplication of the monad $!$ on \mathbf{Cat} (we have $\mu_{\mathbf{C}^{op}} : !!\mathbf{C}^{op} \rightarrow !\mathbf{C}^{op}$, which we can view as a functor from $??\mathbf{C} = (!!\mathbf{C}^{op})^{op}$ to $?\mathbf{C} = (!\mathbf{C}^{op})^{op}$).

Finally, we can compose $\Phi : ?\mathbf{C} \dashrightarrow \mathbf{C}'$ and $\Psi : ?\mathbf{C}' \dashrightarrow \mathbf{C}''$: $\Psi \bullet \Phi = \Psi \circ \Phi^\flat$. Explicitly, $(\Psi \bullet \Phi)(\vec{C}, C'')$ is given by the following formula:

$$\int^{C'_1, \dots, C'_p} \Psi(\vec{C}', C'') \times \int^{\vec{D}_1, \dots, \vec{D}_p} \Phi \vec{D}_1 C'_1 \times \dots \times \Phi \vec{D}_p C'_p \times ?\mathbf{C}[(D_1^1, \dots, D_p^{i_p}), \vec{C}]$$

As promised, we recover Kelly's (Section 4) and Joyal's (Section 5) settings as instances:

1. When $\mathbf{C} = \mathbf{C}' = \mathbf{C}'' = \mathbf{1}$, then Ψ and Φ are presheaves Y, X , and the definition boils down to $Y \bullet X$.
2. When $\mathbf{C} = \mathbf{0}$ (the initial category) and $\mathbf{C}' = \mathbf{C}'' = \mathbf{1}$, then Ψ is a presheaf X , Φ is a set z , and the definition boils down to $\text{Lan}_{\subseteq} Xz$.

It is also easily checked that the identity I as defined in Section 4 is indeed the identity of $\mathbf{Prof}_?$.

We end the section by giving a formula for $\Psi \bullet \Phi$ when $!$ is now an *arbitrary* monad $(!, \eta!, \mu!)$ over \mathbf{Cat} given together with a distributive law: $\lambda : ! \circ Psh \rightarrow Psh \circ !$. This could open the way for handling variations on the shapes of operations (the third kind of variation considered in the introduction) in a profunctor setting.

Recall (Section 6) that given $\Phi : \mathbf{C} \dashrightarrow \mathbf{C}'$ we write Φ' for its presentation as a functor from \mathbf{C}'^{op} to $\mathbf{Set}^{\mathbf{C}}$. Then we can give abstract versions of the steps (1)^{op}, (2)^{op} and (3) above, as follows: (we still write $?\mathbf{C} = (!\mathbf{C}^{op})^{op}$):

- (1)^{op} Given $\Phi : \mathbf{C} \dashrightarrow \mathbf{C}'$, we can define $?\Phi : ?\mathbf{C} \dashrightarrow ?\mathbf{C}'$ by the equation:

$$(?\Phi)' = \lambda \circ !(\Phi')$$

where on the right hand side we apply $!$ as a functor on \mathbf{Cat} .

- (2)^{op} The comultiplication $\delta : ?\mathbf{C} \dashrightarrow ??\mathbf{C}$ is defined by the formula

$$\delta' = \eta_{Psh} \circ (\mu!)_{\mathbf{C}^{op}}$$

- (3) The composition in \mathbf{Prof} of $\Phi' : \mathbf{C}'^{op} \rightarrow \mathbf{Set}^{\mathbf{C}}$ and $\Psi' : \mathbf{C}''^{op} \rightarrow \mathbf{Set}^{\mathbf{C}'}$ is defined as $\Phi'^{\#} \circ \Psi'$. Hence, given $\Phi : ?\mathbf{C} \dashrightarrow \mathbf{C}'$, we obtain the following formula for $\Phi^\flat : ?\mathbf{C} \dashrightarrow ?\mathbf{C}'$:

$$\Phi^{\flat'} = (\eta_{Psh} \circ (\mu!)_{\mathbf{C}^{op}})^{\#} \circ \lambda \circ !(\Phi') = Psh((\mu!)_{\mathbf{C}^{op}}) \circ \lambda \circ !(\Phi')$$

Finally, expanding the definition of Psh , we get the following formula for the composition $\Psi \bullet \Phi = \Psi \circ \Phi^\flat$ in $\mathbf{Prof}_?$:

$$(\Psi \bullet \Phi)(\gamma, C'') = \int^{\gamma'} \Psi(\gamma', C'') \times \int^{\delta} \lambda((! \Phi') \gamma') \delta \times ? \mathbf{C}[\mu_! \delta, \gamma]$$

Following the style of [[26]], one may also hide the distributive law and write

$$(\Psi \bullet \Phi)(\gamma, C'') = \int^{\gamma'} \Psi(\gamma', C'') \times \overline{\gamma'}(\Phi') \gamma$$

where $\overline{\gamma'}(\Phi') \gamma = \hat{\Phi}(\gamma, \gamma')$ can be described abstractly as follows. For each (pseudo)-!-algebra $(\mathbf{D}, \alpha : !\mathbf{D} \rightarrow \mathbf{D})$, for each category \mathbf{C} , and each object C of $!\mathbf{C}$, C induces a functor $\overline{C} = \mathbf{D}^{\mathbf{C}} \rightarrow \mathbf{D}$ defined as $(F \mapsto \alpha(!FC))$. The above formula is obtained by instantiating \mathbf{C} , C , \mathbf{D} , and α as \mathbf{C}'^{op} , γ' , $\mathbf{Set}^{!(\mathbf{C}'^{op})}$, and $Psh(\mu_!) \circ \lambda$, respectively. That this is a (pseudo)-!-algebra structure is established using the equalities satisfied by λ .

This leads us to a generalised definition of operad.

Definition 11.1 *Given a monad $!$ on \mathbf{Cat} and a distributive law $\lambda : ! \circ Psh \rightarrow Psh \circ !$, a \mathbf{C} -coloured !-operad is a monoid in $\mathbf{Prof}_?[\mathbf{C}, \mathbf{C}]$, that is, a functor $X : ?\mathbf{C} \times \mathbf{C}^{op} \rightarrow \mathbf{Set}$ given together with two natural transformations $e : id \rightarrow X$ and $m : X \bullet X \rightarrow X$ satisfying the monoid laws. A $\mathbf{1}$ -coloured !-operad is called a !-operad.*

Hence the non-symmetric operads (resp. symmetric operads, clones) are the $\mathbf{1}$ -coloured $!_{\emptyset}$ -operads (resp. $!_{\{\sigma\}}$ -operads, $!_{\{\sigma, \delta, \epsilon\}}$ -operads), and the non-symmetric coloured operads (resp. symmetric coloured operads) are the coloured $!_{\emptyset}$ -operads (resp. coloured $!_{\{\sigma\}}$ -operads). Indeed, coloured operads were defined in this way by Dolan and Baez [[1]] (though they do not explicate the underlying distributive law or lifting, cf. Footnote 5).

12 Cooperads and properads

So far, we have addressed the variation on the “first axis” that goes from non-symmetric operads to clones. In this section, we address the variation on the second axis (cooperations, bioperations).

Cooperations are dual to operations. Sets of cooperations are organised into cooperads, the notion dual to that of operad. One works now in the Kleisli category $\mathbf{Prof}_!$, i.e., a cooperad is a profunctor $X : \mathbf{1} \multimap !\mathbf{1}$ with a monoid structure in $\mathbf{Prof}_!$, and we can vary on the first axis as we did for operads.

For bioperations, the natural idea is to consider profunctors from $?\mathbf{1}$ to $!\mathbf{1}$. Such profunctors can compose provided there is a distributive law (another one!) $\lambda : ?_X !_X \rightarrow !_X ?_X$: given $\Phi, \Psi : ?\mathbf{1} \multimap !\mathbf{1}$, we define $\Psi \bullet \Phi$ by composing the comultiplication of $?$, $?\Phi$, λ , $!\Psi$, and the multiplication of $!$. The identity is the composition of the counit of $?$ and of the unit of $!$.

This idea has been carried out in detail by Garner, in the case of $X = \{\sigma\}$ [[17]]. We set

$$\lambda((m_1, \dots, m_p), (n_1, \dots, n_q)) \neq \emptyset \quad \text{iff} \quad m_1 + \dots + m_p = n_1 + \dots + n_q$$

and when the equality holds, $\lambda((m_1, \dots, m_p), (n_1, \dots, n_q))$ is the set of permutations s from $m_1 + \dots + m_p$ to $n_1 + \dots + n_q$ such that the graph

- whose set of vertices is the disjoint union of

$$\{1, \dots, p\}, \{1, \dots, m_1 + \dots + m_p\}, \{1, \dots, n_1 + \dots + n_q\}, \text{ and } \{1, \dots, q\}$$

- and whose set of edges is the union

- of the edges between 1 and 1, \dots , m_1 , between 2 and $m_1 + 1, \dots, m_2$, \dots , and between p and $m_{p-1} + 1, \dots, m_p$,
- of the edges between i and $s(i)$ ($i \leq m_1 + \dots + m_p$), and
- of the edges between 1, \dots , n_1 and 1, \dots , and between $n_{q-1} + 1, \dots, n_q$ and q .

is connected. This formalises the idea that when composing two sets of bioperations, we are interested primarily in the compositions that preserve connectedness. More precisely, composition of properations has both a vertical aspect and a horizontal aspect. The distributive law takes care of the vertical aspect. Disjoint connected compositions can be placed in parallel and composed horizontally. Thus, a full categorical account requires a double category setting, that takes care of these two aspects. We refer to [[17]] for details.

Monoids for this composition operation provide an analogue of operads for bioperations, and for $X = \{\sigma\}$ and λ as described above, what we obtain (replacing **Set** by **Vect**) is exactly the notion of *properad* introduced by Vallette in his Thèse de Doctorat [[27, 28]].

Acknowledgements. I collected the material presented here for an invited talk at the conference Operads 2006. When preparing this talk, I benefited a lot from discussions with Marcelo Fiore and Martin Hyland. The string diagrams have been drawn with **strid**, a tool due to Samuel Mimram and Nicolas Tabareau (<http://strid.sourceforge.net/>). I also wish to thank the anonymous referee for his helpful remarks. He in particular pointed out an alternative way to address the size issue, namely to restrict attention to presheaves that are limits of small diagrams of representables.

References

- [1] J. Baez and J. Dolan, Higher-dimensional algebra III: n -categories and the algebra of opetopes, *Advances in Mathematics* 135, 145206 (1998).
- [2] M. Barr and C. Wells, *Toposes, triples and theories*, Springer-Verlag (1985).

- [3] J. Beck, Distributive laws, *Lecture Notes in Mathematics* 80, 119-140 (1969).
- [4] J. Bénabou, Introduction to bicategories, *Lecture Notes in Mathematics* 40, 1-77 (1967).
- [5] J. Bénabou, Les distributeurs, *Université Catholique de Louvain, Institut de Mathématique Pure et Appliquée, Rapport* 33 (1973).
- [6] F. Borceux, *Handbook of categorical algebra*, vol. I: basic category theory, Cambridge University Press (1994).
- [7] F. Borceux, *Handbook of categorical algebra*, vol. II: categories and structures, Cambridge University Press (1994).
- [8] A. Burroni, T-catégories (catégories dans un triple), *Cahiers de Topologie et Géométrie Différentielle* XII(3), 215-321 (1971).
- [9] A. Burroni, Higher Dimensional Word Problem, *Theoretical Computer Science* 115, 43-62 (1993).
- [10] E. Cheng, M. Hyland, J. Power, Pseudo-distributive laws, *Electronic Notes in Theoretical Computer Science* 83 (2004).
- [11] E. Cheng, Weak n-categories: opetopic and multitopic foundations, *Journal of Pure and Applied Algebra* 186, 109137 (2004).
- [12] B.J. Day, On closed categories of functors, *Reports of the Midwest Category Seminar IV, Lecture Notes in Mathematics* 137, 1-38 (1970).
- [13] M. Fiore, Mathematical models of computational and combinatorial structures. Invited address for Foundations of Software Science and Computation Structures (FOSSACS 2005), *Lecture Notes in Computer Science* 3441, 25-46 (2005).
- [14] M. Fiore, N. Gambino, M. Hyland, and G. Winskel, The cartesian closed bicategory of generalised species of structures, *Journal of the London Mathematical Society* 77 (2), 203-220 (2008).
- [15] M. Fiore, N. Gambino, M. Hyland, and G. Winskel, Kleisli Bicategories, in preparation.
- [16] N. Gambino, On the coherence conditions for pseudo-distributive laws, preprint, *ArXiv:0907:1359* (2009).
- [17] R. Garner, Polycategories, PhD Thesis, Cambridge University (2005).
- [18] J.-Y. Girard, Linear logic, *Theoretical Computer Science* 50, 1-102 (1987).
- [19] G.B. Im and G.M. Kelly, A universal property of the convolution monoidal structure, *Journal of Pure and Applied Algebra* 43, 75-88 (1986).

- [20] A. Joyal, Foncteurs analytiques et espèces de structures, in Combinatoire énumérative, Lecture Notes in Mathematics 1234, 126-159 (1986).
- [21] M. Kelly, On the operads of J.P. May, Reprints in Theory and Applications of Categories 13, 1-13 (2005).
- [22] Y. Lafont, Towards an algebraic theory of Boolean circuits, Journal of Pure and Applied Algebra 184 (2-3), 257-310, Elsevier (2003).
- [23] T. Leinster, Higher operads, higher categories, London Mathematical Society Lecture Note Series 298, Cambridge University Press (2004).
- [24] S. Mac Lane, Categories for the working mathematician, Springer-Verlag (1971).
- [25] F. Marmolejo, Distributive laws for pseudomonads, Theory and Applications of Categories 5, 91-147 (1999).
- [26] M. Tanaka and J. Power, Pseudo-distributive laws and axiomatics for variable binding, Higher-Order and Symbolic Computation 19(2-3), 305 - 337 (2006).
- [27] B. Vallette, Dualité de Koszul des PROPS, Thèse de Doctorat, Université de Strasbourg (2003).
- [28] B. Vallette, A Koszul duality for props, Trans. Amer. Math. Soc. 359, 4865-4943 (2007).